

Alessandro Garassino\* & Giovanni Pasini\*\*

Studies on Permo-Trias of Madagascar. 5.  
*Ambilobeia karojoi* n. gen., n. sp. (Crustacea,  
Decapoda) from the Lower Triassic (Olenekian) of  
Ambilobé region (NW Madagascar)

**Abstract** - The study of *Ambilobeia* n. gen., n. sp. (infraorder Penaeidea de Haan, 1849, family Penaeidae Rafinesque, 1815) increases the knowledge about the rich and important faunal assemblage of decapod crustaceans of the Lower Triassic (Olenekian) of the Ambilobé region (NW Madagascar), already widely analysed by Garassino and Teruzzi (1995). The study specimen, recently collected during a surveying, was discovered in a new outcrop located near the village of Bobasatrana. Some morphological features, such as the very elongate rostrum, lacking supra- and subrostral teeth, the presence of one strong tooth located at the base of rostrum, and pereiopods I-III with short and thin chelae, distinguish this new genus from *Ifasya* Garassino and Teruzzi, 1995.

**Key words:** Crustacea, Decapoda, Lower Triassic, NW Madagascar.

**Riassunto** - Studi sul Permo-Trias del Madagascar. 5. *Ambilobeia karojoi* n. gen., n. sp. (Crustacea, Decapoda) del Triassico inferiore (Olenekiano) della regione di Ambilobé (Madagascar nordoccidentale).

Lo studio del nuovo genere *Ambilobeia* (infraordine Penaeidea de Haan, 1849, famiglia Penaeidae Rafinesque, 1815) arricchisce le conoscenze dell'importante fauna a crostacei decapodi del Triassico inferiore (Olenekiano) della Regione di Ambilobé, già ampiamente analizzata da Garassino and Teruzzi (1995). L'esemplare esaminato, raccolto in una recente prospezione, è stato rinvenuto in un nuovo affioramento localizzato in prossimità del villaggio di Bobasatrana. Alcuni caratteri morfologici, quali il rostro estremamente allungato, privo di denti sopra- e sottorostrali, la presenza di un dente robusto localizzato alla base del rostro e i pereiopodi I-III provvisti di chele corte e sottili, discostano questo nuovo genere da *Ifasya* Garassino and Teruzzi, 1995.

**Parole chiave:** Crustacea, Decapoda, Triassico inferiore, NO Madagascar.

## Introduction

The fossiliferous levels of the Lower Triassic marine sediments located S-SW of the village of Ambilobé (Fig. 1), about 150 km SW of Diego Suarez (Antsiranana) have been known since the beginning of last century (Besairie, 1932).

\* Museo Civico di Storia Naturale, C.so Venezia 55, 20121 Milano, Italia. e-mail: a.garassino@tin.it

\*\* Museo Civico dei Fossili di Besano, Via Prestini 5, 21050 Besano (Varese), Italia. e-mail: fossilia@freeonline.zzn.com



Fig. 1 – The point indicates the Bobasatrana outcrop (Ambilobé region, NW Madagascar) where the study specimen was discovered.

Fig. 1 – Il punto indica il giacimento di Bobasatrana (regione di Ambilobé, NO Madagascar) dal quale proviene l'esemplare studiato.

This paper deals with the description of a new genus of decapod crustacean of the Lower Triassic of the Ambilobé region (NW Madagascar), increasing our knowledge about this decapod faunal assemblage, that has been intensively studied by Garassino and Teruzzi (1995). The fossiliferous rocks are included within a sedimentary succession of Lower Triassic age, forming a discontinuous band, about 120 km long, oriented SW-NE. The band of marine sedimentary rocks lies within a depression between two cuestas, formed by the prepalaeozoic crystalline basement to the NE, and by non-fossiliferous continental Middle and Upper Triassic sandstones (Isalo) to the SW.

According to Besairie (1932, 1972), the Lower Triassic succession in the region includes the following (from bottom to top):

- 1 – Neopermian (*Productus*, *Spirifer* and *Xenaspis* beds)
- 2 – Neopermian (*Cyclolobus* and *Xenaspis* beds)
- 3 – Lower Eotrias (*Claraia* and fish beds)
- 4 – Middle Eotrias (fish and ammonite beds)
- 5 – Upper Eotrias (Iraro shales at W and Barabanja *Flemingites* beds at E)
- 6 – Isalo continental sandstones.



Fig. 2 – Bobasatrana outcrop is located near the Mahavavy river.

Fig. 2 – Il giacimento di Bobasatrana è situato in prossimità del fiume Mahavavy.

The thickness of the various members of the series is highly variable (Besairie, 1972). Permian strata are estimated to be about 100 m thick, the marine Lower Triassic series varies from 600 m in the western parts of the basin, in the Ankitokazo area, to 1.200 m in the Andavakoera Massif region, and to just 300 m in the Barabanja basin, on the eastern margin of the sedimentary band.

Beltan (1996) proposed, for the fossil-bearing beds of the Lower Triassic of the Ambilobé region, a Dienerian age (Lower Triassic), after a correlation with the *Lystrosaurus* and *Cyuoguathus* zones of the Beaufort Group (Battail *et al.*, 1987).

Recently, Yanbin *et al.* (2002) studied some conchostracans of the Ambilobé region, ascribing them to *Euestheria (Maguestheria) truemipyi* (Kozur and Seidel, 1983). Since this species was previously discovered also in Bernburg Formation (lowermost Olenekian) of the German basin, we can date the faunal assemblage of Ambilobé region not to Induan (Dienerian), but to Olenekian.

The famous faunal assemblage comprising invertebrates (ammonites, nautiloids, bivalves, annelids, decapod crustaceans, thylacocephalans, cycloids, limulids and conchostracans) and vertebrates (mostly fishes and rare amphibians) is found in the most western outcrops of the band of sedimentary rocks, roughly between the small village of Anaborano Ifasy in the SW and Bobasatrana in the NE (Fig. 1). The first discoveries of fossils in the region was during the fifties of the 20<sup>th</sup> century, during geological prospecting carried out by Bureau de Géologie du Madagascar to the search of gold deposits. The first scientific paper was by Douvillé (1910). Since then many studies were carried out on the invertebrate fauna, mainly dealing with ammonites and bivalves, but also with nautiloids, crustaceans (decapods and thylacocephalans), and polychaete annelids (Vaillant-Couturier Treat, 1933; Van Straelen, 1933; Collignon, 1933, 1934; Alessandrello, 1990; Arduini, 1990). Recently, fossil bivalves from the Lower Triassic of Ambilobé region, identified as *Bakevelliella* sp. cfr. *B. stockleyi* (Cox) (Fang Zhongjie, com. pers.), were found associated with the conchostracan species *Euestheria (Maguestheria) trueiupyi* (Kozur and Seidel, 1983). This bivalve was previously discovered in the marine intercalated beds of the upper part of Ruhembe bed in Kidodi area and the Tanganyka and Sakamena beds of SW Madagascar. In Madagascar, this level lies somewhat below the horizon bearing the amphibian *Rhinesuchus*, which is an important element of the Early Triassic *Lystrosaurus* zone (Beltan, 1996). Therefore, the bivalve-bearing bed is of Early Triassic age. The most recent studies about invertebrates are by Yanbin *et al.* (2002) on conchostracans, by Brambilla *et al.* (in press) on the first record of cycloids in Gondwana, and by Hauschke *et al.* (in press) on the limulids. We point out that the presence of conchostracans, typical organisms of freshwater environments, questions the marine origin of these basins, argued by past and recent studies about the Lower Triassic fauna of Ambilobé region. Also, vertebrates were the subject of many studies, concerning fishes (Priem, 1924), Moy-Thomas (1935), Piveteau (1927, 1934, 1946), Lehman (1948, 1952, 1953, 1956), Beltan (1968, 1996), Olsen (1984) and Barbieri (1991), and amphibians (Lehman, 1955, 1958, 1960, 1963, 1966; Piveteau, 1936, 1938, 1956).

The present research is part of a program of studies started in the eighties of the last century by the Department of the Invertebrate Palaeontology of Museo Civico di Storia Naturale di Milano on the collection of the Lower Triassic fossils of the Ambilobé region. This research program was carried out by the co-operation of the late Dr. Jeannot Rasoanaivo, Director of Service de la Géologie du Madagascar. The specimen subject of this study was collected by one of the authors (G. Pasini).

## Previous studies of decapod crustaceans from Ambilobé region

The first description of decapod crustaceans from the Lower Triassic of Madagascar was that of Van Straelen (1933) who described two incomplete specimens from the Ambilobé region, housed in the collection of the Museum National d'Histoire Naturelle of Paris, ascribing them to *Antrimpos madagascariensis*. Garassino & Teruzzi (1995) enlarged the study of decapod crustaceans of Ambilobé region thanks to a sample of 400 specimens, partly collected during a field mission in 1989 organised by the Department of the Invertebrate Palaeontology of Museo Civico di Storia Naturale di Milano and partly collected in previous years by G. Pasini himself. This research program was allowed by the precious co-operation of the Service de la Géologie du Madagascar. This resulted in recognition of *Ifasya* Garassino and Teruzzi, 1995, with the two species *I. madagascariensis* (Van Straelen, 1933), and *I. straeleni* Garassino and Teruzzi, 1995.

## Material

The present study is based on one specimen, discovered in a new outcrop surrounding the village of Bobasatrana (Figs. 1, 2). The specimen is preserved more or less flattened inside a subellipsoidal nodule, 1.5 cm thick, and 6.5 cm long. It consists of the external mould and preserves also the remains of the internal structures, such as the mandible and the thoracic somites.

Acronym: MSNM: Museo Civico di Storia Naturale di Milano.

### Systematic Palaeontology

Infraorder Penaeidea de Haan, 1849

Superfamily Penaeoidea Rafinesque, 1815

Family Penaeidae Rafinesque, 1815

Genus *Ambilobeia* nov.

*Diagnosis:* subrectangular carapace; long rostrum without supra- and subrostral teeth; one strong tooth at the base of the rostrum; cervical spine and hepatic groove present; pereiopods I-III with chelae short and stout; somite VI strongly elongate.

*Type species:* *Ambilobeia karojoi* n. sp.

*Etymology:* the trivial name alludes to Ambilobé village around which the fossiliferous outcrops are located.

*Description:* same as for the type species.

*Ambilobeia karojoi* n. sp.

Figs. 3, 4

*Diagnosis:* same as for the genus.

*Etymology:* the trivial name alludes to Mr. Karojo, a local prospector who donated the study specimen.

*Holotype:* MSNM i25459.

*Type locality:* Bobasatrana (Ambilobé region); the study specimen comes from a new outcrop surrounding the village of Bobasatrana, discovered recently by




Fig. 3 – *Ambilobeia karojoi* n. gen., n. sp., reconstruction (reconstruction of distal extremity of rostrum, tail fan and pleopods are hypothetical).

Fig. 3 - *Ambilobeia karojoi* n. gen., n. sp., ricostruzione (la ricostruzione dell'estremità distale del rostro, del ventaglio caudale e dei pleopodi è ipotetica).

deforestation to obtain new agricultural lands, and different from the typical outcrop, reported by French palaeontologists in the second half of the last century.

*Geological age:* Olenekian (Lower Triassic).

*Occurrence and measurements:* we ascribe to this species one specimen (MSNM i25459).

MSNM i25459 : total length of the body: 8 cm

length of rostrum: 1 cm

length of carapace excluding rostrum: 2.5 cm

width of carapace: 1.5 cm

length of abdomen: 4 cm

length of somite VI: 1 cm

*Description.* Carapace subrectangular in lateral view (Fig. 3) with ventral margin rising slightly in anterior third. Dorsal margin straight. Rostrum strongly elongate without supra- and subrostral teeth. One strong tooth at base of rostrum. Posterior margin, slightly convex in lower third, strengthened by a thin marginal carina. Ventral margin curvilinear. Ocular incision narrow and shallow. Weak antennal and pterygostomial angles. Dorsal surface of carapace with cervical spine and hepatic groove. Abdominal somites I-V subrectangular, increasing in length posteriorly. Posterior margin of somites I-III straight, while that of somite IV-V posteriorly projecting. Somite VI subrectangular and longer than others. Tail fan incomplete. Eye-stalk short. Scaphocerite with a large lam-




Fig. 4 - *Ambilobeia karojoi* n. gen., n. sp., holotype, n. cat. MSNM i25459 (x1.9).  
Fig. 4 - *Ambilobeia karojoi* n. gen., n. sp., olotipo, n. cat. MSNM i25459 (x1.9).

ina rounded at distal extremity, bearing a spine on outer margin. Pereiopods I-III chelate, increasing in length posteriorly. Chelae short and stout. Pleopods not preserved.

*Discussion.* Garassino and Teruzzi (1995) described the new genus *Ifasya* from Ambilobé region which included *I. straeleni* Garassino and Teruzzi, 1995, and *I. madagascariensis* (Van Straelen, 1933). *Ifasya* Garassino and Teruzzi, 1995 exhibits the following morphological features: subrectangular carapace, long rostrum with or without subrostral median tooth, strong hump at the base of rostrum, deep hepatic groove, weak branchiocardiac groove, gastro-orbital, hepatic and branchiocardiac spines, pereiopods I-III with strong and elongate

chelae, well developed propodus of pereiopod III, strongly elongate somite VI, and triangular telson with pointed distal extremity.

The study of this new specimen pointed out some morphological features, such as the elongate rostrum without supra- and subrostral teeth, strong tooth at the base of the rostrum, pereiopods I-III with short and stout chelae, that are different from species of *Ifasya* Garassino and Teruzzi, 1995. Therefore, we justify the institution of this new genus.

Only the Upper Triassic genus *Autrimpos* Münster, 1839 could show some morphological affinities in the shape of rostrum and body with *Aubilobeia* n. gen. However, the rostrum with supra- and subrostral teeth, the presence of one strong antennal spine, and the presence of hepatic, orbito-antennal and gastro-frontal grooves distinguishes *Autrimpos* Münster, 1839 from the new genus.

### Acknowledgements

We wish to thank Mr. Karojo, a local prospector who donated us the study specimen, R.M. Feldmann, Geology Department, Kent State University, and G. Teruzzi, Department of the Invertebrate Palaeontology of Museo Civico di Storia Naturale di Milano for the useful advice in the draft of manuscript, careful review and criticism of manuscript. Photographic materials by L. Spezia, line drawing by F. Fogliazza.

### References

- Alessandrello A., 1990 – Studies on Permo-Trias of Madagascar. 2. *Palaeoaphrodite auaboraoensis*, a new species of polychaete annelid from the Scythian of Madagascar. *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milaue*, Milano, 131 (10): 205-208.
- Arduini P., 1990 – Studies on Permo-Trias of Madagascar. 1. Thylacocephala from the Lower Trias of Madagascar. *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milaue*, 131 (9): 197-204.
- Barbieri L., 1991 – Poissons Eotriasiques du Nord-Ouest de Madagascar: Systematique et palaeoecologie. *Mémoire de D.E.A. de Paléontologie*.
- Battail B., Beltan L. & Dutuit J.-M., 1987 – Africa and Madagascar during Permo-Triassic time: the evidence of the Vertebrate faunas. In: Gondwana Six, Columbus, Ohio (U.S.A.). *Geophys. Monogr.*, Washington, 41: 147-155.
- Beltan L., 1968 – La faune ichthyologique de l'Eotrias du NW de Madagascar. Le Neurocrane. *Cahiers de Paléontologie*, CNRS, Paris.
- Beltan L., 1996 – Overview of systematics, paleobiology and paleoecology of Triassic fishes of northwestern Madagascar. Mesozoic Fishes. Systematics and Paleoecology. G. Arratia & G. Viohl (eds.). *Verlag Dr. F. Pfeil*, München.
- Besairie H., 1932 – Sur le Permo-Trias marin du nord de Madagascar et l'age du Karroo. *Crit. Soc. Géol. Fr.*, Paris, 10.
- Besairie H., 1972 – Géologie de Madagascar. 1. Les terrains sédimentaires. *Ann. Géol. de Madagascar*, Antananarivo, 35.
- Brambilla S., Garassino A., Pasini G. & Teruzzi G., 2002 – Studies on Permo-Trias of Madagascar. 6. First record of cycloidea from the Lower Triassic

- (Olenekian) of Ambilobé region (NW Madagascar). *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milano*, Milano, 143 (I): 105-115.
- Collignon M., 1933 – Le Trias inférieur marin du nord de Madagascar et sa faune de Céphalopodes. *Cr. Ac. Sc.*, Paris, 197 (18).
- Collignon M., 1934 – Les Céphalopodes du Trias inférieur de Madagascar. *Ann. Paléont.*, Paris, 22 (3-4); 23 (1).
- Douvillé H., 1910 – Sur la découverte du Trias marin à Madagascar. *Bull. Soc. Géol. Fr.*, Paris, 10 (4): 124-133.
- Garassino A. & Teruzzi G., 1995 – Studies on Permo-Trias of Madagascar. 3. The decapod crustaceans of the Albilobé region (NW Madagascar). *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milano*, Milano, 134 (1): 85-113.
- Hauschke N., Wilde V. & Brauckmann C., in prep. – Triassic limulids from Madagascar – missing links in the distribution of Mesozoic Limulacea.
- Lehman J.-P., 1948 – Sur la présence du genre *Birgeria* (Paléoniscidae) dans l'Eotrias de Madagascar. *Cr. Ac. Sc.*, Paris, 226: 426-428.
- Lehman J.-P., 1952 – Etude complémentaire des Poissons de l'Eotrias de Madagascar. *Kunge. Vetensk. Ak. Handt* 1952, ser. 4, 2 (6).
- Lehman J.-P., 1953 – Etude d'un Perleides du Trias de Madagascar. *Ann. Paléont.*, Paris, 39.
- Lehman J.-P., 1955 – Les Stégocéphales du nord-est de Madagascar. *Ann. Paléont.*, Paris, 241: 82-86.
- Lehman J.-P., 1956 – Compléments à l'étude des genres *Ecrinesomus* et *Bobasatrancia* de l'Eotrias de Madagascar. *Ann. Paléont.*, Paris, 42.
- Lehman J.-P., 1958 – Les Vertébrés triasiques inférieurs de Madagascar. *Cr. Réunion CCTA*, Tananarive.
- Lehman J.-P., 1960 – Les Stégocéphales du Trias de Madagascar. *Ann. Paléont.*, Paris, 47.
- Lehman J.-P., 1963 – Les Stégocéphales de Madagascar. *Anu. Géol. Mad.*, Paris, 33.
- Lehman J.-P., 1966 – Nouveaux Stégocéphales de Madagascar. *Ann. Paléont.*, Paris, 2 (2): 117-139.
- Moy-Thomas J.L., 1935 – The Coelacanth fishes from Madagascar. *Géol. Mag.*, Paris, 72 (851).
- Olsen P.E., 1984 – The skull and pectoral girdle of the parasemionotid fish *Watsonulus eugnathoides* from the Early Triassic Sakamena Group of Madagascar, with comments on the relationship of the holostean fishes. *J. Ver. Paleont.*, California, 4 (3): 481-499.
- Piveteau J., 1927 – Sur quelques Poissons fossiles du nord de Madagascar. *Cr. Ac. Sc.*, Paris, 184 (21): 1264-1266.
- Piveteau J., 1934 – Les Poissons du Trias inférieur. Les Actynoptérigiens. *Ann. Paléont.*, Paris, 23 (3-4): 83-178.
- Piveteau J., 1936 – Une forme ancstrale des Amphibiens anoures dans le Trias inférieur de Madagascar. *Cr. Ac. Sc.*, Paris, 202 (19): 1607-1608.
- Piveteau J., 1938 – Un amphibien du Trias inférieur. Essai sur l'origine et l'évolution des Amphibiens anoures. *Ann. Paléont.*, Paris, 26 (3-4): 135-177.
- Piveteau J., 1946 – Les Poissons du Trias inférieur. La famille de Saurichthydés. *Ann. Paléont.*, Paris, 31.
- Piveteau J., 1956 – Présence d'Amphibien Labyrinthodont dans le Trias inférieur du nord de Madagascar. *CRS. Soc. Géol. Fr.*, Paris, 14.

- Priem F., 1924 – Les Poissons fossiles. *Ann. Paléont.*, Paris, 12.
- Vaillant-Couturier Treat I., 1933 – Le Permo-Trias de Madagascar. *Ann. Paléont.*, Paris, 22: 39-95.
- Van Straelen V., 1933 – *Antrimpos madagascariensis* Crustacé décapode de Permotrias de Madagascar. *Bull. Mus. roy. Hist. nat. Belg.*, Bruxelles, 9 (15): 1-3.
- Yanbin S., Garassino A. & Teruzzi G., 2002 – Studies on Permo-Trias of Madagascar. 4. Early Triassic conchostracans from Madagascar. *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milano*, Milano, 143 (I): 3-11.

Ricevuto: 2 aprile 2002

Approvato: 2 maggio 2002